


1. Multiple Choice: Electric Field of Discrete Charges: 

## Question

1) A charge  $q_1 = 10\text{nC}$  is placed on the positive  $y$ -axis at  $\vec{r}_1 = (0, 6\text{m}, 0)$ , and another charge  $q_2 = 10\text{nC}$  is placed on the negative  $y$ -axis at  $\vec{r}_2 = (0, -6\text{m}, 0)$ . The  $x$ -component of the electric field at  $\vec{r} = (8\text{m}, 0, 0)$  is :



## Answer

 3.4 N/C 2.3 N/C 1.4 N/C 0.32 N/C 0.023 N/C 0.0082 N/C

2. Multiple Choice: Pb Nucleus in E Field:

Question

2) A fully ionized Pb nucleus (mass  $m = 3.5 * 10^{-25}$  kg, charge  $+82 e$ ) is released from rest in a uniform electric field (ignore gravity). After 5 seconds, the nucleus is moving at a speed of  $2 * 10^6$  m/s. The magnitude of the electric field is :



Answer

 2.1 V/m 1.2 V/m 0.68 V/m 0.35 V/m 0.011 V/m 0.0012 V/m

3. Multiple Choice: Electric Flux: 

## Question

3) You hold a plate of area  $A$  inside a uniform electric field such that the normal vector of its plane makes an angle of 30 degrees with the field lines. You measure an electric flux of  $25 \text{ Vm}$  through the plate. You then rotate the plate such that it is still fully immersed in the field but its normal vector is parallel to the field lines. The new flux is :

Flux =  $25 \text{ Vm}$ 

after rotation Flux = ?


## Answer

 50 Vm 29 Vm 25 Vm 22 Vm 12.5 Vm

4. Multiple Choice: Charge inside Conducting Shell: 

## Question

4) A point charge  $Q$  is placed at the center of a conducting spherical shell with inner radius  $R_1$  and outer radius  $R_2$ . The total charge on the shell is  $-3Q$ . The charge on the outer surface of the shell (at  $r = R_2$ ) is equal to :



## Answer

 -3Q -2Q -Q 0 +Q +2Q +3Q



5. Multiple Choice: Insulating Sphere:

Question

5) A solid insulating sphere of radius  $R$  is uniformly charged with total charge  $Q$  and placed at the origin. Use Gauss' law to determine the electric field for a radius  $r < R$ . The answer is :

Answer

$$E(r) = kQ/r^2$$


$$E(r) = kQ/(2 r^2)$$

$$E(r) = kQ/(R r)$$

$$E(r) = kQr / (2 R^3)$$

$$E(r) = kQr/R^3$$

$$E(r) = kQr^2/(2 R^4)$$

6. Multiple Choice: Circular Ring: 

Question

6) A thin circular ring of radius  $R$  and uniformly distributed charge  $Q$  is placed in the  $x$ - $y$ -plane, with its center at the origin. The electric potential of the ring along the  $z$ -axis is given by  $V(z) = kQ/\sqrt{z^2 + R^2}$ . The electric field on the  $z$ -axis is given by :

Answer

$$k Q \operatorname{sign}(z) / (z^2 + R^2)$$

$$2 k Q \operatorname{sign}(z) / (z^2 + R^2)$$

$$\checkmark k Q z / (z^2 + R^2)^{3/2}$$

$$- k Q z / (z^2 + R^2)^{3/2}$$

$$k Q z^2 / (z^2 + R^2)^2$$

$$2 k Q z^2 / (z^2 + R^2)^2$$

$$k Q z^3 / (z^2 + R^2)^{3/2}$$


$$- k Q z^3 / (z^2 + R^2)^{3/2}$$

## 7. Multiple Choice: Work:

## Question

7) A configuration of electric field lines (solid lines) and corresponding equipotential surfaces (dashed lines) is shown on the right.

The work done by the electric force when a positive charge  $q$  is moved from A to B,  $W_{AB}$ , is related to the work done for the same charge to move from C to D,  $W_{CD}$ , as :



## Answer

$$W_{AB} > W_{CD}$$

$$W_{AB} = W_{CD}$$

$$W_{AB} < W_{CD}$$


not uniquely determined.

8. Multiple Choice: Potentials: 

## Question

8) A configuration of electric field lines (solid lines) and corresponding equipotential surfaces (dashed lines) is shown on the right.

Comparing the potentials at points C and D, one has :



## Answer

$V_D > V_C$

$V_D = V_C$

$V_D < V_C$

not uniquely determined.



9. Multiple Choice: Equilibrium:

Points: 7

Question

9) In electrostatic equilibrium, the electric potential inside a conductor is :

Answer

always zero.

always a constant.

increases from the negatively charged surface to the positively charged surface.

decreases from the negatively charged surface to the positively charged surface.

10. Multiple Choice: Two Charges: 

Question

10) A positive charge  $q_1=3\text{nC}$  is located at the origin, and an unknown charge  $q_2$  at  $x_2= 0.16\text{m}$ . The electric potential is found to be zero at  $x=0.12\text{m}$ . The value of  $q_2$  is :


Answer

 -4 nC -3 nC -2 nC -1 nC 1 nC 2 nC 3 nC 4 nC

## 11. Multiple Choice: Charged Disk:

## Question

11) A thin circular disk of radius  $R$  has a surface charge density of  $\sigma = B r^2$  with  $B$  given. The potential of the disk at its center (that is, at the origin) is given by :



## Answer

$$V = k \pi B \frac{R^2}{2}$$

$$V = k \pi B R^2$$

$$V = 2 k \pi B R^2$$

$$V = k \pi B R^3$$

$$V = k \pi B \frac{R^3}{2}$$

$$V = 2 k \pi B \frac{R^3}{3}$$

## 12. Multiple Choice: Charge in Field:

Question

12) An electron (charge  $q = -e$ ) is placed in a static electric field at a point  $P_0$  where the electric potential is  $+12\text{ V}$ . The electron is released from rest. It then reaches a point  $P_1$  where it is observed to have a kinetic energy of  $4.2 \times 10^{-18}\text{ J}$ . Calculate the electric potential at  $P_1$ .

Answer

$V_1 = -48\text{ V}$

$V_1 = -29\text{ V}$

$V_1 = -13\text{ V}$

$V_1 = -0.5\text{ V}$

$V_1 = 16\text{ V}$

$V_1 = 38\text{ V}$

$V_1 = 74\text{ V}$



13. Multiple Choice: Energy of System of Charges: 

Question

13) Three equal negative point charges,  $-Q$ , are placed on the corners of an equilateral triangle with sides of equal length  $D$ . The electric potential energy stored in this system is :

Answer

$$-6 k Q^2 / D$$

$$-3 k Q^2 / D$$

$$-2 k Q^2 / D$$

$$-k Q^2 / D$$

$$k Q^2 / D$$

$$2 k Q^2 / D$$

$3 k Q^2 / D$

$$6 k Q^2 / D$$