



1. [Qid 12] (4 points) There are two arcs of charge that each create  $3/4$  of a circle and are centered on the origin as shown below. Both arcs have the same uniform charge density  $\lambda$  where  $\lambda$  is a positive value. What direction will the net electric field point at the origin? (All angles are given as counterclockwise from the positive  $x$ -axis.)

- (A)  $135^\circ$
- (B)  $-\hat{x}$
- (C)  $+\hat{x}$
- (D)  $+\hat{y}$
- (E)  $45^\circ$
- (F)  $225^\circ$
- (G)  $-\hat{y}$
- (H)  $315^\circ$



2. [Qid 3] (8 points) A point charge  $Q = -500$  nC and two unknown point charges,  $q_1$  and  $q_2$ , are placed as shown in the figure where all positions are given in units of meters. The electric field at the origin  $O$ , due to charges  $Q$ ,  $q_1$  and  $q_2$ , is equal to zero. The charge  $q_1$  is closest to

- (A) 141 nC
- (B) -244 nC
- (C) 244 nC
- (D) -141 nC
- (E) 281 nC
- (F) -281 nC
- (G) -315 nC
- (H) 315 nC




3. [Qid 5] (6 points) Three protons exist at the corners of an equilateral triangle of side length 2 cm. If two protons are held in place and the third is released from rest, what maximum speed will the third proton reach?

- (A) 5.26 m/s
- (B) 36.7 m/s
- (C) 7.30 m/s
- (D) 14.2 m/s
- (E) 27.6 m/s
- (F) 3.72 m/s

4. [Qid 8] (10 points) There is an electric field  $\vec{E} = 3.00\hat{i} - 2.00y\hat{j}$  N/C in the region that includes a cube of side length  $L = 10.0$  m and is oriented as shown in the figure below. What is the net charge inside the cube?


- (A) -8.85 nC
- (B) +3.42 nC
- (C) +8.85 nC
- (D) +17.7 nC
- (E) -3.42 nC
- (F) -17.7 nC



5. [*Qid 11*] (6 points) An object is comprised of 2,500,000 total particles. The particles can be either protons or electrons. The total charge of this object is  $-1.44 \times 10^{-13}$  C. How many of these particles are protons?
- (A) 600,000
  - (B) 800,000
  - (C) 900,000
  - (D) 1,900,000
  - (E) 2,500,000
  - (F) 1,300,000
  - (G) 0
  - (H) 1,700,000


6. [Qid 1] (6 points) In the Chapter 21 Part 2 recitation, we found that for an insulating ring of charge with radius  $a$  and charge density  $\lambda(\theta) = \lambda_0 \sin \theta$  where  $\theta$  is measured counterclockwise from the positive  $x$ -axis, the electric field at the center was  $\vec{E} = -\frac{k\lambda_0\pi}{a}\hat{j}$ . What is the sign of the potential at the center of this ring due to that charge distribution?

- (A) Negative
- (B) Zero
- (C) Positive




7. [Qid 10] (10 points) A very long insulating cylinder with a radius  $R = 4.00$  cm is uniformly charged with a volume charge density  $\rho = 5.00$  nC/m<sup>3</sup>. Find the magnitude of an electric field at the distance  $r = R/2$  from the axis of the cylinder.
- (A) 11.3 N/C
  - (B) 565 N/C
  - (C) 30.2 N/C
  - (D) 5.65 N/C
  - (E) 3.77 N/C
  - (F) 3.41 N/C
  - (G) 1130 N/C
  - (H) 341 N/C

8. [Qid 15] (4 points) A metal sphere with a radius  $R$  initially has a charge of  $-Q$ . A second metal sphere with a radius  $2R$  initially has a charge of  $+9Q$ . When the switch below is closed, the two spheres will be connected by a conducting wire. What will happen when this switch is closed?
- (A) The charges will not redistribute at all and will stay where they are.
  - (B) The charge will redistribute until there is the same magnitude of electric field just outside the surface of the spheres.
  - (C) The charge will redistribute until there is the same charge on both spheres.
  - (D) The charge will redistribute until there is the same potential at the surface of both spheres.




9. [Qid 14] (6 points) In the diagram below there are two very large, thin insulating sheets that are separated by a distance  $d$ . You know that the bottom sheet has a surface charge density of  $+5\sigma$ . You also know that the electric field at a distance  $d$  above the top plate has a magnitude of  $E = \frac{2\sigma}{\epsilon_0}$  and is pointing down towards the plates. What is the surface charge density on the top plate?



- (A)  $+3\sigma$   
 (B)  $-1\sigma$   
 (C)  $+9\sigma$   
 (D)  $-3\sigma$   
 (E)  $-7\sigma$   
 (F)  $-9\sigma$   
 (G)  $+1\sigma$   
 (H)  $+7\sigma$
10. [Qid 7] (8 points) A uniform electric field of magnitude 450 N/C is directed in the negative  $y$ -direction. Point A is located at (0.25m, -0.5 m) and point B is located at (0.75m, 0.4m). What is the potential difference  $V_B - V_A$ ?
- (A) -405 V  
 (B) 315 V  
 (C) -513 V  
 (D) 221 V  
 (E) -221 V  
 (F) 405 V  
 (G) 513 V  
 (H) -315 V


11. [Qid 13] (6 points) In the diagram below there is a thick-walled, conducting cubic shell that has total charge of  $-5 \text{ mC}$ . At the center of that is a thick-walled, conducting spherical shell that has a total charge of  $+4 \text{ mC}$ . At the center of that is a point charge. If there is  $+7 \text{ mC}$  on the OUTSIDE surface of the cube, what is the charge of the point charge?

- (A)  $+8 \text{ mC}$
- (B)  $-8 \text{ mC}$
- (C)  $+5 \text{ mC}$
- (D)  $-5 \text{ mC}$
- (E)  $+16 \text{ mC}$
- (F)  $+1 \text{ mC}$
- (G)  $-1 \text{ mC}$
- (H)  $-16 \text{ mC}$



12. [Qid 9] (4 points) A point charge with magnitude  $Q$  is situated in the centrum of two imaginary hemispheres of radii  $a$  and  $b$  as shown below. Compare the fluxes  $\Phi_a$  and  $\Phi_b$  due to this point charge if you know the charge is negative.

- (A)  $\Phi_a > \Phi_b$ ,  $\Phi_a$  is positive and  $\Phi_b$  is negative
- (B)  $\Phi_a < \Phi_b$ ,  $\Phi_a$  is negative and  $\Phi_b$  is positive
- (C)  $\Phi_a = \Phi_b$ , both fluxes are positive
- (D)  $\Phi_a = \Phi_b$ , both fluxes are negative
- (E)  $\Phi_a > \Phi_b$ , both fluxes are positive
- (F)  $\Phi_a < \Phi_b$ , both fluxes are negative






13. [Qid 4] (8 points) In a particular region of space, a 300 mC charge experiences the effects of a potential energy function given by  $U(x, y, z) = 4x^2y + 3xyz - 5y^2z^3$ . What is the electric field vector at the point (3,2,-1)? Assume all values are in SI units.

- (A)  $-108\hat{i} - 9\hat{j} + 72\hat{k}$
- (B)  $-42\hat{i} - 47\hat{j} + 42\hat{k}$
- (C)  $-30\hat{i} + 51\hat{j} + 78\hat{k}$
- (D)  $-72\hat{i} + 18\hat{j} + 20\hat{k}$
- (E)  $-360\hat{i} - 30\hat{j} + 240\hat{k}$
- (F)  $-240\hat{i} + 60\hat{j} - 66.7\hat{k}$
- (G)  $-140\hat{i} - 157\hat{j} + 140\hat{k}$
- (H)  $-100\hat{i} + 170\hat{j} + 260\hat{k}$

14. [Qid 6] (4 points) The figure below shows a series of equipotential surfaces. At which point does the electric field have the greatest magnitude?

- (A) Point A
- (B) Point B
- (C) Point C
- (D) Point D
- (E) Point E



15. [Qid 2] (10 points) A *non-uniform* linear charge characterized by the charge per unit length  $\lambda(y)$  is located on the  $y$ -axis extending from  $y = -a$  to  $y = +a$ . Which of the following integrals give the electric field  $\vec{E}$  on the  $x$ -axis at  $x = +a$ ? Note that  $\lambda(y)$  is not a constant value.

(A)  $\vec{E} = \int_{-a}^{+a} \frac{k\lambda(y)dy}{a^2} \hat{i}$

(B)  $\vec{E} = \int_{-a}^{+a} \frac{ka\lambda(y)dy}{(y^2 + a^2)^{3/2}} \hat{i} - \int_{-a}^{+a} \frac{ky\lambda(y)dy}{(y^2 + a^2)^{3/2}} \hat{j}$

(C)  $\vec{E} = \int_{-a}^{+a} \frac{k\lambda(y)dy}{(y^2 + a^2)^2} \hat{i} + \int_{-a}^{+a} \frac{ky^2\lambda(y)dy}{(y^2 + a^2)^{3/2}} \hat{j}$

(D)  $\vec{E} = 0$

(E)  $\vec{E} = \lambda(y) \int_{-a}^{+a} \frac{kady}{(y^2 + a^2)^{3/2}} \hat{i} - \lambda(y) \int_{-a}^{+a} \frac{kydy}{(y^2 + a^2)^{3/2}} \hat{j}$

(F)  $\vec{E} = \lambda(y) \int_{-a}^{+a} \frac{kdy}{(y^2 + a^2)} \hat{i}$

(G)  $\vec{E} = \lambda(y) \int_{-a}^{+a} \frac{kdy}{(y^2 + a^2)^2} \hat{i} + \lambda(y) \int_{-a}^{+a} \frac{ky^2dy}{(y^2 + a^2)^{3/2}} \hat{j}$

(H)  $\vec{E} = \int_{-a}^{+a} \frac{ka\lambda(y)dy}{(y^2 + a^2)^{3/2}} \hat{i}$

